arXiv:hep-ph/0209039v1 4 Sep 2002

Dual coherent particle emission as generalised Cherenkov-like effect in high energy particle collisions

D. B. Ion[*]
National Institute for Physics and Nuclear Engineering,
NIPNE-HH, Bucharest, P.O. Box MG-6, Romania
and
TH Division, CERN, CH-1211 Geneva 23, Switzerland

E. K. Sarkisyan[†]
EP Division, CERN, CH-1211 Geneva 23, Switzerland
and
High Energy Physics Institute Nijmegen (HEFIN),
University of Nijmegen/NIKHEF,
NL-6525 ED Nijmegen, The Netherlands

(Dated: February 1, 2008)

In this paper we introduce a new kind of nuclear/hadronic coherent particle production mechanism in high-energy collisions called *dual coherent particle emission (DCPE)* which takes place when the phase velocities of the emitted particle v_{Mph} and that of particle source v_{B_1ph} satisfy the dual coherence condition: $v_{Mph} \leq v_{B_1ph}^{-1}$. The general signatures of the DCPE in the nuclear and hadronic media are established and some experimental evidences are given.

PACS numbers: 25.40.-h, 25.70.-z, 25.75.-q, 13.85.-t

Cherenkov radiation was first observed in the early 1900's by the experiments developed by Mary and Pierre Curie when studying radioactivity emission. The nature of such radiation was unknown until the experimental works (1934-1937) of P.A. Cherenkov and the theoretical interpretation by I. E. Tamm and I. M. Frank (1937) when it was clarified that this radiation is produced by charged particles travelling through matter at speeds larger than the phase velocity of light in the medium. Although it was not well understood and recognised until 1958 (Nobel Prize in Physics), the effect afterwards was called the Cherenkov radiation (CR). The idea that meson production in nuclear interaction may be described as a process similar to the electromagnetic CR go back to Wada, Ivanenko and Gurgenidze, and Blohintzev and Indenbom (see Refs. [1]) who propagated it first on a qualitative basis and after that by using some nuclear models. In the last decade this idea of the generalisation of the CR to mesons and photons inside the nuclear and hadronic media was systematically investigated theoretically [2-4] and also tested experimentally [5-8]. Such idea was also extended to a Cherenkov-like mechanism for meson production in hadron-hadron interactions [8-11] at high energies. In this analogy, a fast hadron is the source of mesonic field as it is the fast charged particle in the normal Cherenkov radiation. Thus the mesonic Cherenkov-like effect might be expected to occur when a *hadron traverses the nuclear (or hadronic) matter with a velocity exceeding the phase velocity of the mesonic field.*

In this paper, we introduce a new kind of coherent particle production mechanism in particle collisions called *dual coherent particle emission (DCPE)* from which all kind of the generalised Cherenkov-like effects can be obtained. The DCPE effects are expected to take place when the phase velocities of the emitted particle v_{Mph} and that of particle source v_{B_1ph} satisfy the *dual coherence condition*: $v_{B_1ph}v_{Mph} \leq 1$. We investigate the general signatures of DCPE effects in nuclear and hadronic media and a propose their experimental tests. The comparison of the absolute predictions [3] on the nuclear pionic Cherenkov-like effect (NPICR) with the recent experimental results obtained in relativistic nuclear collisions [5-7] is given.

Dual coherent particle emission (DCPE). Let us start with a general $\mathrm{B}_1 \rightarrow \mathrm{MB}_2$ decay described in Fig. 1(a). Here a particle M [with energy ω, momentum $k = \mathrm{Re}n_M(\omega)\sqrt{\omega^2 - M_{\mathrm{M}}^2}$, rest mass M_{M}, and refractive index $n_M(\omega)$] is emitted in a (nuclear, hadronic, dielectric, etc.) medium from an incident particle B_1 [with energy E_1, momentum $p_1 = \mathrm{Re}n(E_1)\sqrt{E_1^2 - M_1^2}$, rest mass M_1, and refractive index $n_1(E_1)$] that itself goes over into a final particle B_2 [with energy E_2, momentum $p_2 = \mathrm{Re}n(E_2)\sqrt{E_2^2 - M_2^2}$, rest mass M_2, and refractive index $n_2(E_2)$].

Here we prove that in order to obtain a genuine spontaneous particle emission in a given medium *the two general conditions* are necessary to be fulfilled:

- (i) The incident particle-source must be coupled to a specific radiation field (RF) (see Fig. 1) and that the particles propagation properties in medium must be modified.
- (ii.1) The particle source must be moving in medium with a dual phase velocity $v_{B_1ph}^{-1}$ higher than the phase velocity v_{Mph} of the RF-quanta.



FIG. 1: (a) The two-body decay process $\mathrm{B}_1 \to \mathrm{MB}_2$ in nuclear media. (b) The nuclear mesonic Cherenkov-like radiation (NMCR) limit [3]: $v_{Mph} \leq v_1$. (c) The nuclear baryonic Cherenkov-like radiation limit [3]: $v_{B_2ph} \leq v_1$

- (ii.2) The particle source must be moving in medium with a dual phase velocity $v^{-1}_{B_1ph}$ higher than v_{B_2ph}.

Proof: The propagation properties of particles in a medium are changed in agreement with their elastic scattering with the constituents of that medium. So, the phase velocity $v_{Xph}(E_X)$ of any particle X (with the total energy E_X and rest mass M_X) in medium is modified as follows:

$$v_{Xph}(E_X) = \frac{E_X}{p_X} = \frac{1}{\mathrm{Re}n_X(E_X)} \cdot \frac{E_X}{[E_X^2 - M_X^2]^{\frac{1}{2}}} \tag{1}$$

The *refractive index* $n_X(E_X)$ in a medium composed from the constituents "c" can be calculated in standard way by using the Foldy-Lax formula [12] (we work in the units system $\hbar = c = 1$)

$$n_X^2(E_X) = 1 + \frac{4\pi\rho}{E_X^2 - M_X^2} \cdot C(E_X)\overline{f}_{Xc\to Xc}(E_X) \tag{2}$$

where ρ is the density of the constituents, $C(\omega)$ is a *coherence factor* [$C(\omega) = 1$ when the the medium constituents are randomly distributed], $\overline{f}_{Xc\to Xc}(E_X)$ is the averaged forward Xc-scattering amplitude.

Now, by using the *energy-momentum conservation law* for the decay $\mathrm{B}_1 \to \mathrm{MB}_2$ in medium,

$$E_1 = E_2 + \omega, \quad \overrightarrow{p}_1 = \overrightarrow{p}_2 + \overrightarrow{k} \tag{3}$$

we obtain (see angles definition in Fig. 1):

$$cos\theta_{1k} = v_{Mph}v_{B_1ph} + \frac{1}{2p_1k}[-D_{B_1} + D_{B_2} - D_M] \tag{4}$$

$$cos\theta_{12} = v_{B_1ph}v_{B_2ph} + \frac{1}{2p_1k}[-D_{B_1} - D_{B_2} + D_M] \tag{5}$$

$$cos\theta_{2k} = v_{Mph}v_{B_2ph} + \frac{1}{2p_2k}[D_{B_1} - D_{B_2} - D_M] \tag{6}$$

where D_X, $X \equiv B_1, B_2, M$, are departures from mass shell, and are given by

$$D_X \equiv E_X^2 - p_X^2 = M_X^2 + [1 - (\mathrm{Re}n_X(E_X))^2][E_X^2 - M_X^2] \tag{7}$$

We note that the second terms in the right side of Eqs (4)-(6) can be considered as quantum corrections to the first "classical" terms [3].

Now, a rigorous proof of the statement (ii) is obtained from conditions that the respective emission angles must be the physical angles. The *coherence quantum conditions: $cos\theta_{ij} \leq 1$, $i, j = 1, 2, k$,* (4)-(6), at the high incident particles energies transform into *classical coherence conditions*, e. g.,

$$cos\theta_{1k} = v_{Mph}v_{B_1ph} \leq 1 \tag{8}$$

which is equivalently to

$$v_{Mph} \leq v^{-1}_{B_1ph}, \text{ or } v_{B_1ph} \leq v^{-1}_{Mph} \tag{9}$$

It is worth to note that from the *dual coherence conditions* (9) we obtain the following DCPE condition.

- (iii) In case when $v^{-1}_{B_1ph} = v_1$, from the condition (ii) the two important generalised Cherenkov-like limits follow: (iii.1) the NMCR Cherenkov-like radiation limit $v_{Mph} \leq v_1$, and (iii.2) the NBCR Cherenkov-like radiation limit $v_{B_2ph} \leq v_1$.

The proof of the statement (iii) is obtained immediately if one observes that when the particle B_1 is *on the mass shell in medium* ($\mathrm{Re}n_1 = 1$) *then* $v^{-1}_{B_1ph}$=v_1, and the *dual coherence conditions* (ii.1) and (ii.2) will go into the *Cherenkov-like coherence conditions* (iii.1) and (iii.2), respectively.

For each kind of generalised Cherenkov-like effect there is a *threshold projectile kinetic energy* given by

$$T_{thr}(E_X) = M_p\sqrt{1 - [v_{Xph}(E_X)]^2 - 1}]. \tag{10}$$

Now, we can obtain a classification of these DCPE effects not only on the basis of four fundamental (strong, electromagnetic, weak, and gravitational) interactions but also using the above *"M-B duality"* as well as the *crossing symmetry.*

The main signatures of the (DCPE) as generalised Cherenkov-like effects are as follows:

- The differential cross sections posses the bumps in the energy bands where the DCPE-coherence conditions are fulfilled.
- The DCPE-effects are threshold mechanisms.

- Coherent particles emitted via the DCPE mechanism must be coplanar with the incoming and outgoing projectiles: strong (θ_{1k}, ω) and (θ_{1k}, E_p) correlations.

- The intensities as well as the absorption effects can be calculated as in the case of generalised Cherenkov-like effects [3] by using Feynman diagrams in medium.

- Any two-body decay process $B_1 \rightarrow MB_2$ in medium via DCPE mechanism posses two limiting modes: the NMCR mode in which the particle M is emitted when the coherence condition $v_{Mph} \leq v_1$is fulfilled, and the NBCR mode when the coherence condition $v_{B_2ph} \leq v_1$ is fulfilled.

COHERENT MESON EMISSION IN NUCLEAR MEDIA. The lightest mesons for which the elementary scattering amplitudes on nucleons in the forward direction are well known from the experimental data are pions [13]. In the last decade, the *nuclear pionic Cherenkov-like radiation (NPICR)* was intensively investigated theoretically [2,3] on the basis of elementary pion-nucleon scattering amplitudes in the forward direction. The characteristic features of the NPICR-pions predicted in Refs. [2,3] are as follows:

- The numerical values on the refractive index of pions in nuclear medium was obtained in the standard way by applying Lax-Foldy formula (2) with the coherence factor $C = 1$.

- The NPICR-coherence condition $v_{ph}(\omega) \leq v$, was found to be *fulfilled in the three energy bands:*
  (CB1) 190 MeV$\leq \omega \leq$ 315 MeV, for all $\pi^{\pm,0}$;
  (CB2) 910 MeV $\leq \omega \leq$ 960 MeV, only for π^+;
  (CB3) $\omega \geq$ 80 GeV, for all $\pi^{\pm,0}$.
  The values for the NPICR-thresholds for all the three pionic bands are shown in Fig. 2.

- The NPICR-pions must be coplanar with the incoming and outgoing projectile possessing strong (θ_{1k}, ω) and (θ_{1k}, T_p) correlations (see Fig. 7 from Ref. [3]).

- The *NPICR-differential cross sections (DCS)* are peaked at the energy ω_m = 244 MeV for CB1-emission band when absorption in medium is taken into account. The CB1-peak width in the DCS is predicted to be $\Gamma_m \leq$ 25 MeV (see details in Figs. 8 and 9 from Ref.[3]).

- The energy behaviour of NPICR-peak position is predicted to be as T_p^{-2} (see Fig. 12 in [3]), while the A-target dependence of the NPICR differential cross sections is given in Fig. 14 of [3].



FIG. 2: The absolute predictions for the threshold $T_{thr}(\omega)$ for single coherent meson production via *pionic Cherenkov-like radiation (NPICR)* mechanism [3] in the nuclear reaction ^{208}Pb (p$\rightarrow$ Nπ).



FIG. 3: The energy distributions obtained [5,6] for the π^-s produced in the central Mg-Mg collisions at 4.3 GeV/c per nucleon.

The existence of the NPICR-CB1-emission band [2,3] was recently confirmed experimentally by the JINR-Dubna group [5,6] in the studies of Mg-Mg central collisions at 4.3 GeV/c per nucleon by processing the pictures from the 2m Streamer Chamber SKM-200. After processing a total number of 14218 events, which were found to meet the centrality criterion, the experimental results shown in Fig.3 have been obtained.

As one sees from Fig. 3, the energy distributions of emitted pions in the central collisions have a significant peak over the inclusive background. The experimental values of the peak position and its width are

$$E^*_m = 238 \pm 3(\mathrm{stat}) \pm 8(\mathrm{syst})\ \mathrm{MeV}, \qquad (11)$$

$$\Gamma_m = 10 \pm 3(\mathrm{stat}) \pm 5(\mathrm{syst})\ \mathrm{MeV}. \qquad (12)$$

Hence, the *values of the peak energy and its width* obtained in Refs. [5,6] *are in a good agreement with the absolute NPICR predictions for* $\omega_m and \Gamma_m$ [2,3]. It is important to note that the above value of E^*_m is similar to the position of the peak observed in Ref. [15] in the analysis of π^+production in coincidence measurements of (p,n) reactions at 0^0 on C, the effect connected with the NPICR mechanism [3].

Therefore, we conclude that the NPICR cross sections in all emission bands, as well as other NPICR-signatures, are large enough in order to be experimentally measured in exclusive experiments. Of the great importance is the experimental discovery of the first coherent NPICR band in the inclusive measurements since it represents the first experimental proof that the generalised Cherenkov-like effect is the real mechanism for the coherent particles production with the rest mass different from zero.

GENERALISED MESONIC CHERENKOV-LIKE RADIATION IN HADRONIC MEDIA. The mesonic Cerenkov-like radiation in hadronic media was considered by many authors [8-11]. A systematic investigation of the classical and quantum theory of this kind of effects in hadronic media can be found in Ref. [9]. The classical variant [2,9] of the Cherenkov mechanism was applied to the study of single meson production in hadron-hadron interactions at high energies. This variant is based on the usual assumption that hadrons are composed from a central core in which most of the hadron mass is concentrated surrounded by a large and more diffuse mesonic cloud (hadronic medium). Then, it was shown [9,11] that a *hadronic mesonic Cherenkov-like radiation (HPICR)* with an mesonic refractive index given by a pole approximation, can be able to explain the integrated cross sections of a single meson production in hadron-hadron interaction. As an example, in Fig. 4 we present the integrated cross sections of the process pp → ppπ^0, compared with the HPICR-predictions in the *limit of the HPICR mechanism dominance*. This result was very encouraging for the extension of HPICR-dominance hypothesis to other single meson production in hadron-hadron collisions at high energies. Collecting all the χ^2/dof for all 139 reaction fitted with the *hadronic mesonic Cherenkov-like* (HMCR) mechanism, we get the surprisingly good description as shown in Fig. 5. We must underline that only reactions with single meson production was fitted (a single parameter fit) with the HMCR predictions on the integrated cross sections.

Also, recently, in hadronic experiments [5-7,16], the Cherenkov-like radiation in the variant proposed in [10], has been observed.



FIG. 4: Experimental statistical test of the *HMCR-mechanism dominance* in hadron-hadron collisions.



FIG. 5: The number of reactions fitted with HMCR-predictions for integrated cross sections as a function of χ^2/dof.

CONCLUSIONS. In this paper a new mechanism called *dual coherent particle emission* (DCPE) is introduced.

The results and conclusion obtained in this paper are as follows:

- The DCPE mechanism includes in a more general and exact way all the types of the Cherenkov mechanisms thanks to the *general dual coherence condition* $v_{B_1ph}v_{Mph} \leq 1$ which has the *particular limiting cases: M-Cherenkov-like radiation limit:* $v_{Mph}\leq v_1$*, and B-Cherenkov-like radiation limit:* $v_{B_2ph} \leq v_1$.

- In the *nuclear media* we presented the first experimental results [5,6] which confirm with high accuracy the predictions [3] in the first CB1-pionic band.

- In the *hadronic media* we presented the *first statistical test* (see Figs. 4 and 5) based on the fit of the integrated cross sections of 139 reaction of single meson production. These results suggest a possible *HMCR-dominance* of the single meson production in high energy hadron-hadron collision.

Finally, it is important to note that special experimental techniques based on the coincidence measurements are necessary in order to extract the yields of the DCPE generalised Cherenkov-like effects from the background produced by other mechanisms.

We would like to thank G. Altarelli for fruitfull discussions. One of the authors (D.B.I.) would like to thank TH Division for hospitality during his stay at CERN. One of the authors (E.K.S.) would like to thank the "Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO)" for support.

---

* Electronic address: Dumitru.Ion@cern.ch
† Electronic address: sedward@mail.cern.ch

[1] W. Wada, Phys. Rev. 75, 981 (1949); D. Ivanenko and V. Gurgenidze, Dokl. Acad. Nauk SSSR 67, 997 (1949); D. I. Blohintzev and V. L. Indenbom, JETF 20, 1123 (1950); V. L. Indenbom, JETF 21, 737 (1951); D. B. Ion, Stud. Cercet. Fiz. 22, 125 (1970).

[2] D.B. Ion and W. Stocker, Phys. Lett. B 273, 20 (1991); Phys. Rev. C48, 1172 (1993); Acta Phys. Pol. B24, 1785 (1993), Phys. Lett. B 346, 172 (1995); Rom. J. Phys. 43, 15 (1998).

[3] D.B. Ion and W. Stocker, Phys. Rev. C 52, 3332 (1995), and references therein.

[4] D. B. Ion and W. Stocker, Phys. Lett. B 258, 262 (1991) and B 262, 498 (1991); Ann. Phys. (N.Y.) 213, 355 (1992); Phys. Lett. B 311, 339 (1993); Phys. Lett. B 323, 446 (1994).

[5] G.L. Gogiberidze, L.K. Gelovani, and E. K. Sarkisyan, Phys. Lett. B 471, 257 (1999).

[6] G. L. Gogiberidze, E. K. Sarkisyan and L. K. Gelovani, Nucl. Phys. B (Proc. Suppl.) 92, 75 (2001).

[7] G. L. Gogiberidze, L.K. Gelovani, and E.K. Sarkisyan, Physics of Atomic Nuclei 64, 143 (2001).

[8] W. Czyz, T. Ericson, and S. L. Glashow, Nucl. Phys. 13, 516 (1959) ; W. Czyz and S. L. Glashow, Nucl. Phys. 20, 309 (1960); G. Yekutieli, Nuovo Cimento 13, 446 (1959); 13, 1306 (1959); P. Smrz, Nucl. Phys. 35, 165 (1962); Czech. J. Phys. B12 (1962) 169.

[9] D. B. Ion, *Mesonic Cherenkov-like effect as possible mechanism of meson production in hadronic interactions,* Doctoral Thesis, Bucharest University, 1971; D. B. Ion and F. G. Nichitiu, Nucl. Phys. B 29 (1971) 547.

[10] I. M. Dremin, JETP Lett. 30, 140 (1979); Sov. J. Part. Nucl. 18, 31 (1987).

[11] D. B. Ion, Rev. Roum. Phys. 36 (1991), 585 and 595; D. B. Ion and F. G. Nichitiu, Rom. J. Phys. 37 (1992) 763; D. B. Ion, A. Rosca and C. Petrascu, Rom. J. Phys 37 (1992) 991; D. B. Ion and A. Rosca 38 (1993) 655; D. B. Ion, A. Rosca and C. Petrascu, Rom. J. Phys 39 (1994) 395.

[12] L. L. Foldy, Phys. Rev. 67 (1945) 107; M. Lax, Rev. Mod. Phys. 23 (1951) 287; H. Feshbach, Ann. Rev. Nucl. Sci. 8 (1958) 49; M. L. Goldberger and K. M. Watson, *Collision Theory*, pp 766-775, Wiley, New York-London-Sydney, 1964.

[13] E. Pedroni et al., Nucl. Phys. A 300 (1978) 321; G. Hohler et al., *Handbook of Pion-Nucleon Scattering,* Karlsruhe Report Nr. 12-1, 1979.

[14] M. Pardy, Phys. Lett. B 336, 362 (1994).

[15] J. Chiba et al., Phys. Rev. Lett. 67, 1982 (1991).

[16] I. M. Dremin et al., Sov. J. Nucl. Phys. 52, 536 (1990); EHS-NA22 Collab., N.M. Agagabyan et al., Phys. Lett. B 389, 397 (1996); Shaoshung Wang et al., Phys. Lett. B 427, 385 (1998).